



CENTRE FOR EVIDENCE-BASED CONSERVATION

**Effectiveness of the Control of Ragwort (*Senecio*) Species
Systematic Review Series:**

SYSTEMATIC REVIEW N^o. 5b.

**“Can biological control by the use of natural enemies effectively
control *Senecio jacobaea* (common ragwort)?”**

REVIEW REPORT

**COMPILED BY
Roberts, P.D. & Pullin, A.S.**

**Centre for Evidence-Based Conservation
School of Biosciences
The University of Birmingham
Edgbaston
Birmingham
B15 2TT
U.K.**

February 2005

This review should be cited as:

Roberts, P.D. & Pullin, A.S. (2005) *Effectiveness of the control of ragwort (*Senecio*) species: “Can biological control by the use of natural enemies effectively control *Senecio jacobaea* (common ragwort)?”* Systematic Review N^o. 5b. Centre for Evidence-Based Conservation, Birmingham, U.K.

COVER SHEET

Title	Can natural enemies effectively control common ragwort (<i>Senecio jacobaea</i>)
Reviewer(s)	Roberts, P.D., (PDR) & Pullin, A.S. (ASP)
Date protocol first published on website	28 th June 2004
Date review first published on website	March 2005
Date of most recent amendment	-
Date of most recent SUBSTANTIVE amendment	-
Most recent changes	-
Contact address	Philip Roberts Centre for Evidence-Based Conservation School of Biosciences University of Birmingham Edgbaston Birmingham B15 2TT UK Tel: 0121 4144090 Email: pdr387@bham.ac.uk
Systematic review N ^o .	N ^o . 5 part b
Sources of support	NERC, English Nature.
Conflict of interests	None reported.

SUMMARY

BACKGROUND

Biological control is the purposeful introduction of natural enemies (biocontrol agents) by land managers and scientists as a means to weaken and/or suppress invading plants or animals. Since the 1930s natural enemies, such as *Tyria jacobaeae* and *Longitarsus jacobaeae* have been used in an attempt to control *Senecio jacobaea* due to its potential to poison livestock and cause economic loss to agriculture. This systematic review uses explicit methodology to capture and evaluate primary evidence for the effectiveness of natural enemies as control agents of *S. jacobaea*.

OBJECTIVE

To assess the effectiveness of *T. jacobaeae* (cinnabar moth); *L. jacobaeae* (ragwort flea-beetle); or a combination of both for the control of *S. jacobaea* (common ragwort).

SEARCH STRATEGY

Electronic databases: ISI Web of Knowledge (WoK) containing ISI Web of Science and ISI Proceedings; Science Direct; JSTOR; Index to Thesis; UMI ProQuest Digital Dissertations; COPAC (incl. British Library); Natural History Museum Library; AGRICOLA and SCIRUS. English Nature (EN); Countryside Council for Wales (CCW) and Scottish Natural Heritage (SNH) publications were all searched online.

Other searches: Additional references not captured by the initial searches, were located via the inspection of all reference lists of studies accepted at full text.

SELECTION CRITERIA

Studies were included if they fulfilled the following selection criteria:

Subject: All studies which focused on *S. jacobaea* (common / tansy ragwort).

Intervention: The use of the natural enemy, *T. jacobaeae* (cinnabar moth); *L. jacobaeae* (ragwort flea-beetle); or a combination of both to control *S. jacobaea*.

Outcome(s): A measure of the ragwort population densities or a measure of an aspect of the plants characteristics (such as dry weight, capitula per plant or seed viability).

Type of study: All primary, quantitative studies and reports with a comparator of an appropriate control were included within formal meta-analysis. In addition time series studies which lacked a comparator were collated and the relative change in the *S. jacobaea* population was calculated.

DATA COLLECTION AND ANALYSIS

Both reviewers assessed study inclusion / exclusion, methodological quality & data extraction. Any discrepancies were resolved by discussion. Information on the population focus, methodology, interventions and outcomes were abstracted from the original studies into a specially designed, pre-tested spreadsheet. Data synthesis using

standardised mean difference (SMD), random effects model meta-analysis was performed by one reviewer, with the results being discussed by both reviewers.

MAIN RESULTS

Meta-analysis: Although the overall density of *S. jacobaea* plants (SMD $d+ = -0.27$; $p = 0.4473$ ns (positive sensitivity analysis) and SMD $d+ = -0.27$; $p = 0.995$ ns (negative sensitivity analysis)) is not significantly reduced by the *T. jacobaeae* treatment, certain plant characteristics are significantly reduced therefore affecting the reproductive ability of *S. jacobaea* plants: capitula per plant (SMD $d+ = -8.71$; sig. = 0.0076** (positive analysis) and SMD $d+ = -7.90$; $p = 0.0455^*$ (negative analysis)) and seeds per plant (SMD $d+ = -693.92$; $p = 0.0174^*$).

Due to the limited sample sizes for the meta-analyses for *L. jacobaeae* and the combined *T. jacobaeae* & *L. jacobaeae* treatment no robust results could be calculated for their effectiveness in controlling *S. jacobaea*.

Time series: Results from the captured time series datasets show considerable variability in the reduction of the *S. jacobaea* using *T. jacobaeae*, with some sites even showing increases in plant densities. Using *L. jacobaeae* all sites showed a considerable decline (mean = 96.5%; range = 93.1% to 99.9%, $n = 2$) in *S. jacobaea* densities. The combination treatment using both *T. jacobaeae* & *L. jacobaeae* again results in considerable decline (mean = 99.53%; range = 98.46% – 100%, $n = 5$) of *S. jacobaea* densities.

REVIEWERS' CONCLUSIONS

The best available evidence suggests that *T. jacobaeae* reduces the reproductive ability of *S. jacobaea* therefore potentially reducing the further spread of the plant. Although densities of *S. jacobaea* were reduced, the result was not statistically significant. From the time series evidence *T. jacobaeae* showed vast variability for its effectiveness in controlling *S. jacobaea* densities.

There were insufficient datasets available to draw any robust conclusions from the meta-analyses for both *L. jacobaeae* and the combination of both natural enemies on *S. jacobaea*. The additional evidence provided by the time series shows that: *L. jacobaeae* caused major reductions of *S. jacobaea* densities and plant characteristics in all datasets. However, it is the use of the combination treatment; of both *T. jacobaeae* & *L. jacobaeae* that shows the greatest potential for the effective control of *S. jacobaea* densities.

The inclusion of time series datasets, without a comparator, within this systematic review allows for further tentative conclusions to be drawn for all three treatments. These results should be treated with caution due to uncertainty of confounding effects and the reduced methodological quality used to obtain the original datasets.

Further randomised control trials (RCTs) with multiple replicates and at least a two year time period are required to investigate the effectiveness of all three treatments on *S. jacobaea* densities and plant characteristics.

CONTENT PAGE

1. BACKGROUND	6
2. OBJECTIVES	7
3. METHODS	7
3.1. Search Strategy for Identification of Studies	7
3.2. Criteria for Inclusion of Studies within Systematic Review	8
3.2.1. <i>Types of Study</i>	9
3.2.2. <i>Population Focus</i>	9
3.2.3. <i>Interventions of Interest</i>	10
3.2.4. <i>Desired Outcome Measures</i>	10
3.3. Study Quality Assessment	10
3.4. Data Extraction	12
3.5. Data Checking	12
3.6. Data Synthesis	12
4. RESULTS	13
4.1. Results of Search	13
4.2. Results of the Study Methodological Quality	14
5. OUTCOME OF THE REVEIW	16
5.1. The Effectiveness of <i>Tyria jacobaeae</i>	16
5.2. The Effectiveness of <i>Longitarsus jacobaeae</i>	16
5.3. The effectiveness of combining <i>T. jacobaeae</i> & <i>L. jacobaeae</i>	16
5.4. Time series datasets	16
6. DISCUSSION	20
7. REVIEWERS' CONCLUSIONS	21
7.1. Implications for Conservation	21
7.2. Implications for Research	21
8. SOURCES OF SUPPORT	22
9. ACKNOWLEDGEMENTS	22
10. REFERENCES	22
10.1. References to Studies Included in this Systematic Review	22
10.2. References to Studies Excluded from this Systematic Review	23
10.3. Additional References	23
Appendix 1	24
Appendix 2	26

Effectiveness of the control of ragwort (*Senecio*) species: Can biological control by the use of natural enemies effectively control *Senecio jacobaea* (common ragwort).

1. BACKGROUND

Biological control is the purposeful introduction of natural enemies (biocontrol agents) by land managers and scientists as a means to weaken and suppress weed and pest species. Natural enemies are used to decrease an invasive plants' competitive advantage over native species and to weaken the invading population by increasing leaf mortality; decreasing plant size; reducing flower and seed production; and/or limiting population expansion (Cameron, 1935).

The use of natural enemies to control *S. jacobaea* was first undertaken in the early 1930s by the introduction of *Tyria jacobaeae* (cinnabar moth, also previously named *Callimorpha jacobaeae* L.) into Australia. Since then the addition of other natural enemies, such as the widely used *Longitarsus jacobaeae* (ragwort flea-beetle); the rarely used *Cochylis atricapitana* (crown boring moth) and *Botanophila seneciella* (ragwort seed fly) have been employed in an attempt to control the spread of *S. jacobaea*.

The first instar larva (caterpillar) of *T. jacobaeae* is yellow with a black head. Later, they become more brightly coloured with black and yellow bands. They feed initially on the leaves, but as the foodplant matures, they move onto the flowers. They however, rarely attack the rosettes of mature plants unless all the flowering plants have been consumed. Numerous studies give detailed description of the life cycle of *T. jacobaeae* (e.g. Cameron, 1935) and the seeking of a biotype that can be acclimatised to the region where control is desired (Schmidl, 1972).

The adults of *L. jacobaeae* feed on the rosettes and the leaves, causing a holed “shot-gun” appearance, while the larvae eat the roots. There have been two different biotypes described which have been introduced to different regions (Frick, 1971, Frick and Johnson 1973).

Apart from the separate use of these two natural enemies, they have been used as a combined treatment (e.g. Hawkes & Johnson, 1976) in an attempt to control *S. jacobaea* from both ends of the plant (above and below ground).

S. jacobaea is a problem in many countries around the world. Biological control has been undertaken to various degrees of success in Australia, New Zealand, Canada, U.S.A. and Europe (James *et al.*, 1992, Schmidl, 1972). However, the Australian dairy industry estimated that this species still causes \$2,428,211/year (£1,011,595/year) in lost milk production and \$434,327/year (£180,941/year) in lost beef production. Based on these figures alone, the annual cost of *S. jacobaea* to Australia would be in excess of \$4.0 million (>£1.6 million).

2. OBJECTIVES

To evaluate the effectiveness of natural enemies used for the control of common ragwort (*S. jacobaea*), by the use of datasets concerning measures of population density or plant characteristics following biological control experiments.

To explore, when possible, the following reasons for heterogeneity within the datasets: 1. the number of natural enemies; 2. the length of follow-up period; 3. the soil type; 4. altitude of sites; 5. weather / climate of the sites and 6. age of plants.

3. METHODS

3.1. Search Strategy for Identification of Studies

The following electronic databases were searched for the identification of a ragwort library of all possible relevant studies for this systematic review. All dates listed below show the years covered by that particular database.

1. **ISI Web of Knowledge (WoK)** using CrossSearch Form involving the searching of the following products:
 - ISI Web of Science (1981 to present).
 - ISI Proceedings (1990 to present).
2. **Science Direct** – Agricultural and Biological Sciences (1823 to present).
3. **JSTOR**.
4. **Index to Theses** (1970 to 2003).
5. **UMI ProQuest Digital Dissertations** (1950s to 2003).
6. **COPAC** – database of the 24 main British and Irish university libraries and the British Library and National Library of Scotland.
7. **UK Natural History Museum Library** (1980 to present + 80% prior).
8. **AGRICOLA** – two databases for the National Agricultural Library of America:
 - Online Public Access Catalogue (books).
 - Journal Article Citation Index (journals).
9. **SCIRUS** – Scientific Search Engine.
10. **Wildlink** – English Nature's Library Catalogue (only available on-site).

The following search terms were used on all the above electronic databases to identify the initial library of all possibly relevant studies. This created an initial general ragwort control library of studies, from which filtering for particular control methods could take place (section 3.2)

1. Ragwort AND Control
2. *Senecio* AND Control
3. Pulling AND (Ragwort OR *Senecio*)
4. Herbicide AND (Ragwort OR *Senecio*)
5. Spraying AND (Ragwort OR *Senecio*)
6. Wiping AND (Ragwort OR *Senecio*)
7. “Spot Treatment” AND (Ragwort OR *Senecio*)

8. Cutting AND (Ragwort OR *Senecio*)
9. Mechanical AND (Ragwort OR *Senecio*)
10. Biological AND Control AND (Ragwort OR *Senecio*)
11. Cinnabar AND Moth
12. *Tyria* AND *jacobaeae*
13. Ragwort AND “flea beetle”
14. *Longitarsus* AND *jacobaeae*

Searches on www.alltheweb.com were constructed using the search terms below. In all cases only the website that was captured by the alltheweb search was assessed, in the order that they appeared from each search, no links were followed from identified sites. The first 50 websites were assessed for their relevance.

1. Ragwort + control
2. Ragwort + control + results
3. *Senecio* + control
4. *Senecio* + control + results

All of the electronic and web searches were initially completed in May 2004 with additional web searches being undertaken in August 2004 & January 2005.

Relevant organisations such as English Nature, Scottish Natural Heritage, Countryside Council for Wales, The National Trust, UK Wildlife Trusts & Agricultural Libraries were contacted and their website publication lists searched for pertinent grey literature or unpublished data.

Attempts were made to contact first authors of included studies, if any queries about the clarification of the reported results, missing data values or further explanation of their findings were required.

3.2. Criteria for Inclusion of Studies within Systematic Review

As briefly mentioned above, the initial library of studies was developed as a general library enabling further filtering for applicability to a particular review. This was generated from the 14 search terms across the 10 electronic databases (with the removal of all duplications) and the web based searches. See figure one, for the number of studies at each step of the assessment for inclusion relevance.

From the general library an initial inclusion criterion for title and abstract assessment was developed. All studies to be included within the systematic review were required to be focused, or partially focused, on any ragwort species and also to contain any intervention which was undertaken to control/reduce the amount of ragwort present on the site/area. All studies which fulfilled this first assessment at title and abstract or those that lacked sufficient detail to make an assessment were placed in a second library for further assessment at full text.

All reference lists of the remaining studies were checked to identify any additional studies missed in the initial search. These were also added to the second library to give the grand total of studies to be viewed at full text.

Full text assessment of these remaining studies was undertaken with the added inclusion criteria that all studies had to contain either: a study comparator (i.e. a treatment plot and a control plot), compare two or more sites (treatment site and control site) or have a measurement of the plants density / characteristics over a time period **without a comparator** (these are referred to from here as “time series”).

A random subset of 20 studies at full text inclusion was independently assessed by a second reviewer. Cohen’s Kappa analysis was performed to test the level of agreement and repeatability between reviewers. In this case the level of agreement was good, with $K = 0.87$. Any disagreements for a studies inclusion / exclusion were resolved through discussion.

After full text assessment the final library of accepted studies were split in relation to the relevant management intervention used to control ragwort species. For inclusion within this systematic review on the effectiveness of natural enemies the following final inclusion criteria had to be fulfilled.

3.2.1. Types of Study

To be eligible for inclusion within meta-analysis, studies had to contain a comparator, of a treatment/control plot. This criterion therefore allows for the inclusion of randomised control trials (RCTs), control trials (CTs) and site comparisons studies (SCSs). These were collated for each natural enemy and combination prior to the data analysis.

Time series data could not be included within the meta-analysis, due to the lack of a comparator component within the experimental design which is required when calculating each study’s individual point estimate and the overall effect size for the intervention in question. All time series studies that have been identified by the search strategy are however presented in tables, so that the data are available for interested parties and for subsequent analysis.

Studies were not rejected due to country of origin or language of publication. All foreign language papers were translated prior to final full text assessment.

3.2.2. Population Focus

All studies which focused on common / tansy ragwort or *S. jacobaea* were considered for inclusion within the systematic review.

3.2.3. Interventions of Interest

Studies which focused on the biological control of *S. jacobaea* using the following natural enemies:

1. Cinnabar Moth larvae (*Tyria jacobaeae*).
2. Ragwort Flea Beetle larvae & adult (*Longitarsus jacobaeae*).
3. A combination of the both the above.

Studies which focused on the plant-insect population dynamics / interaction were rejected as the main concern of these studies is not normally the eradication / control of *S. jacobaea*.

3.2.4. Desired Outcome Measures

The outcomes of interest for this review were:

- i. Measures of the ragwort population densities.
- ii. Measures of the plant characteristics (e.g. dry weight or seed viability).

Due to the biennial nature of ragwort, it would be ideal for the outcome measures (follow-up period) to cover at least two seasons after the intervention. However, it is anticipated that there would be substantial variation in the length of follow-up, therefore all studies which conformed to the above criteria were accepted and an assessment of the follow-up periods for all datasets would be used as one of the measures to test for heterogeneity within analyses.

3.3. Study Quality Assessment

The quality of each of the accepted papers was assessed in accordance with the study quality assessment instrument (Appendix 1) by one reviewer. The quality assessment involved looking at each individual study design; baseline comparators; intra-treatment variation; measurement of co-/intervention and outcome measures. The details of the study quality assessments were recorded for each study on their individual study characteristic table (see Appendix 2).




Figure 1: A diagrammatic representation of the study selection phase for inclusion of evidence within the systematic process. The number of studies at each stage is represented by N.

3.4. Data Extraction

Information on the population focus; methodology; interventions; and outcomes were abstracted from all the original studies viewed at full text assessment onto a specially designed spreadsheet. Studies which fulfilled all inclusion criteria had the following additional data and methodological characteristics extracted onto the spreadsheet, to allow investigation of instances of heterogeneity within meta-analyses:

Title, year of publication, site location, habitat, experimental area, species of ragwort, age of the plants, soil type, natural enemies used, month application, year of application, number applied, altitude of site, notable previous interventions undertaken, methods of application, time of follow-up, additional applications, other site activities during experiment, number of replicates, method of recording (use of scales, health scores etc.), outcome measure with related standard deviation values (e.g. % data, ragwort plants m², ragwort plants per plot etc.) and adverse effects.

In all cases the mean, number of replicates and the standard deviation measures were required from both the treatment and control to allow meta-analyses to be performed on the datasets. The data that were extracted from the original studies and used within the meta-analyses are presented within the study characteristic tables (Appendix 2). All data were derived from the field/plot level avoiding pseudo-replication.

3.5. Data Checking

Attempts were made to contact first authors of included studies for missing data and standard deviation measures and / or if there were any errors and inconsistencies in the data.

3.6. Data Synthesis

The spreadsheets of extracted data were grouped by natural enemy and used with the StatsDirect™ programme for data synthesis. All outcomes were in the form of continuous data, which were pooled across trials using Standardised Mean Difference (SMD) meta-analysis and the random effects model (Sutton *et al.* 2000).

Sensitivity analyses were performed on the data to determine the effect of the inclusion of non-independent datasets. These are defined as those studies which in their experimental design compared a number of different treatments plots against only one control plot. For non-independent data, effect sizes were generated for all data and combined in meta-analyses of most positive and negative independent effect sizes (sensitivity analyses) to assess the impact of the intervention.

4. RESULTS

4.1. Results of Search

Lists of all studies viewed at each stage of the systematic review study inclusion process are available on request. Of the 14 studies containing evidence on the control of *S. jacobaea* with natural enemies, 9 studies had data / datasets which fulfilled the inclusion criteria for meta-analysis and 8 of the studies contained data / datasets which were time series; these are tabulated in section 5.4 to show trends and for future analysis.

All included studies were published or reporting on experiments conducted between 1953 and 1992. The studies were conducted in: U.S.A. (n = 6), Canada (n = 3), Australia (n = 2), United Kingdom (n = 2) and Netherlands (n = 1).

The natural enemies listed in Tables 1 and 2 had datasets captured by the search strategy. All measures of control (i.e. reduction in density/plant characteristics) were considered for meta-analysis if there were two or more datasets available for synthesis.

Table 1: The source of the datasets which related to each measure of control for *S. jacobaea* populations that were synthesised in meta-analysis for the effectiveness of the natural enemies: a) *Tyria jacobaeae*; b) *Longitarsus jacobaeae*; c) A combination of the two.

A) *Tyria jacobaeae*

	Density of plants	Dry weight (g) per plot	Capitula per plant	Seeds per capitula	Seed per plant	Viability of seeds (%)	Height of plants	Leaves per plant
Bornemissza, G. F. (1966)		Y	Y	Y	Y	Y		
Cameron, E. (1935)					Y			
Ganeshan, S. (1992)	Y		Y				Y	
Harris, P. (1974)	Y							
Harris, P., Thompson, L. S., et al. (1976)	Y							Y
James, R. R., et al. (1992)	Y		Y	Y	Y	Y		Y
Schmidl, L. (1972)		Y	Y			Y	Y	

B) *Longitarsus jacobaeae*

	Density of plants	Capitula per plant	Biomass (g) per plot
James, R. R., et al. (1992)	Y	Y	Y
McEvoy, P. et al. (1991)	Y	Y	Y

C) Combination of *Tyria jacobaeae* & *Longitarsus jacobaeae*

	Density of plants
Hawkes, R. B. & Johnson, G. R. (1976) C	Y
James, R. R., et al. (1992) C	Y

Table 2: The number of datasets (grouped by outcome measure and natural enemy), available for inclusion within meta-analysis. First number = only independent datasets; Number in brackets = all datasets (independent & non-independent data).

	Tyria jacobaeae	Longitarsus jacobaeae	Combination
Density of plants	5(10)	2	3
Dry weight (g) per plot	2(3)	-	-
Capitula per plant	5(8)	2	-
Seed per capitula	2	-	-
Seed per plant	4	-	-
Viability of seeds (%)	3	-	-
Height of plants	3(5)	-	-
Leaves per plant	2(5)	-	-
Biomass (g) per plot	-	2	-

4.2. Results of the Studies Methodological Quality Assessment

Low scores were assigned to the following datasets: Cameron, E. (1935) dataset C; Harris, P., Thompson, L. S., *et al.* (1976); Hawkes, R. B. & Johnson, G. R. (1976); M^cEvoy, P. B. (1985); M^cEvoy, P. *et al.* (1991) datasets B & C; Nagel, W. P. & Isaacson, D. L. (1974); Pemberton, R. W. & Turner, C. E. (1990) and Windig, J. J. (1993) as they were all time series, lacking comparator, information about methodology and experimental area. The highest scores were assigned to: James, R. R., *et al.* (1992) and M^cEvoy, P. *et al.* (1991) dataset A, as they were both randomised controlled trials with potential reasons for heterogeneity clearly stated.

A summary of the study quality assessment is presented overleaf (Table 3) for more details see the individual study characteristic tables (Appendix 2).

Table 3: Summary of the study methodology quality assessment (in descending order) for each of the **datasets** contained in the studies accepted at full text. Please note: studies normally contain more than one dataset as identified within the second column “dataset ID”. For more details of each study see Appendix 2.

Study	Dataset ID	Study design	Country of origin	Baseline comparison	Intra treatment variation	Study quality score
James, R. R., <i>et al.</i> (1992)	A-C	RCT	USA	Size of plots, habitat type, location and age of stand are all homogenous	Stand age, habitat type and location are all homogenous	91
McEvoy, P. <i>et al.</i> (1991)	A	RCT	USA	Size of plots, habitat type, location and age of stand are all homogenous	Stand age, habitat type and location are all homogenous	91
Bornemissza, G. F. (1966)	A-F	CT	Australia	Size of plots, habitat type, location, altitude, age of stand and soil type are homogenous	Altitude, stand age, habitat type, and location are all homogenous	76
Ganeshan, S. (1992)	A-B	CT	UK	Size of plots, habitat type, location and age of stand are all homogenous	Stand age, habitat type and location are all homogenous	70
Hawkes, R. B. & Johnson, G. R. (1976)	B-C	CT	USA	Size of plots, habitat type, location and age of stand are all homogenous	Stand age, habitat type and location are all homogenous	70
Schmidl, L. (1972)	A	CT	Australia	Habitat type, location, and age of stand are all homogenous	Stand age, habitat type and location are all homogenous	69
Cameron, E. (1935)	A-B	HCT	UK	Size of plots, habitat type, location and age of stand are all homogenous	Stand age, habitat type and location are all homogenous	63
Harris, P., Thompson, L. S., <i>et al.</i> (1976)	A-D	SCS	Canada	Size of plots, habitat type, location and age of stand are all homogenous	Stand age, habitat type and location are all homogenous	50
Harris, P. (1974)	A	SCS	UK	Habitat type, location and age of stand are all homogenous	Stand age, habitat type and location are all homogenous	48
Harris, P., Wilkinson, A. T. S. <i>et al.</i> (1976)	B-C	Time series	Canada	Size of plots, habitat type, location, age of stand and soil type are all homogenous	Stand age, habitat type and location are all homogenous	41
Nagel, W. P. & Isaacson, D. L. (1974)	A-D	Time series	USA	Habitat type, location, altitude and age of stand are all homogenous	Altitude, stand age, habitat type and location are all homogenous	41
Windig, J. J. (1993)	A-C	Time series	Netherlands	Size of plots, habitat type, location, soil type and age of stand are all homogenous	Stand age, habitat type and location are all homogenous	41
Harris, P., Wilkinson, A. T. S. <i>et al.</i> (1976)	A	Time series	Canada	Size of plots, habitat type, location and age of stand are all homogenous	Stand age, habitat type and location are all homogenous	40
Hawkes, R. B. & Johnson, G. R. (1976)	A	Time series	USA	Size of plots, habitat type, location and age of stand are all homogenous	Stand age, habitat type and location are all homogenous	40
McEvoy, P. <i>et al.</i> (1991)	B-C	Time series	USA	Size of plots, habitat type, location and age of stand are all homogenous	Stand age, habitat type and location are all homogenous	40
Pemberton, R. W. & Turner, C. E. (1990)	A-C	Time series	USA	Size of plots, habitat type, location and age of stand are all homogenous	Stand age, habitat type and location are all homogenous	40
Cameron, E. (1935)	C	Time series	UK	Size of plots, habitat type, location and age of stand are all homogenous	Stand age, habitat type and location are all homogenous	39
McEvoy, P. B. (1985)	A-B	Time series	USA	Size of plots, habitat type, location and age of stand are all homogenous	Stand age, habitat type and location are all homogenous	39

5. OUTCOME OF THE REVIEW

Studies were sorted according to which natural enemy they studied and the outcome measure used for measuring their effectiveness in controlling *S. jacobaea* populations. DerSimonian-Laird pooled d^+ values give overall pooled effect sizes for each of the meta-analyses (Egger *et al*, 2003; Sutton *et al*, 2000). Negative values for the effect sizes indicate a reduction in the parameter of control being measured for the effectiveness of the natural enemy in reducing *S. jacobaea*. The approximate 95% confidence intervals and DerSimonian-Laird chi squared significance values are also presented for each meta-analysis (Table 4, 5 & 6).

When non-independent data is present, sensitivity analyses were performed. This is when the most positive results (those that had the greatest effect from each dataset) or most negative results (those that had the least effect from each dataset) are re-analysed together thus forming independent datasets to derive the overall effect size (Sutton *et al*, 2000).

It was not possible to investigate reasons for heterogeneity within the meta-analyses due to insufficient recording of the control measures and the limited number of datasets.

5.1. The effectiveness of *T. jacobaeae*

Although the overall density of *S. jacobaea* plants (SMD d^+ = -0.27; p = 0.4473 ns (positive sensitivity analysis) and SMD d^+ = -0.27; p = 0.995 ns (negative sensitivity analysis)) is not significantly reduced by *T. jacobaeae*, certain characteristics are significantly reduced therefore affecting the reproductive ability of *S. jacobaea* (Table 4): capitula per plant (SMD d^+ = -8.71; sig. = 0.0076** (positive analysis) and SMD d^+ = -7.90; p = 0.0455* (negative analysis)) and seeds per plant (SMD d^+ = -693.92; p = 0.0174*). Other plant characteristics of *S. jacobaea* with significant reductions include: seeds per capitula and dry weight of plants, however sample sizes are very small (n = 2).

5.2. The effectiveness of *L. jacobaeae*

The analyses of the three outcome measures show reductions in all characteristics of *S. jacobaea* (Table 5). However, due to the limited number of dataset (sample sizes = 2 for all), none proved significant, as the 95% confidence intervals of each outcome measure crossed the zero line (line of no effect): Density of plants (p = 0.3377); capitula per plant (p = 0.1929) and biomass (g) per plant (p = 0.7736).

5.3. The effectiveness of combining *T. jacobaeae* & *L. jacobaeae*

The analysis of the density of plants showed a very marginal reduction in *S. jacobaea*, but again proved not significant with the confidence intervals crossing the line of no effect: Density of plants (SMD = -0.086; 95% CI = -1.47 to 1.30; p = 0.9028). However, again sample sizes are very small (n = 2).

Table 4: The effectiveness of *Tyria jacobaeae* on the reduction of *S. jacobaea*.
n = number of datasets; CI = confidence intervals and χ^2 sign. = chi-squared significance.

Control Measure	Result		Independent	Positive Sensitivity Analysis	Negative Sensitivity Analysis
Density of plants	<i>Not significant, robust sample size</i>	n Effect Size 95% CI χ^2 sign.		5 -0.27 -0.97 to 0.43 0.4473 (ns)	5 0.002 -0.70 to 0.71 0.995 (ns)
Capitula per plant	<i>Significant, robust sample size</i>	n Effect Size 95% CI χ^2 sign.		5 -8.71 -15.10 to -2.31 0.0076**	5 -7.90 -15.65 to -0.16 0.0455*
Seeds per plant	<i>Significant, robust sample size</i>	n Effect Size 95% CI χ^2 sign.	4 -693.92 -1265.75 to -122.09 0.0174*	Not required	Not required
Viability (%) of seeds	<i>Very significant, robust but small sample size</i>	n Effect Size 95% CI χ^2 sign.	3 -0.74 -1.11 to -0.36 < 0.0001***	Not required	Not required
Seeds per capitula	<i>Very significant, robust but very small sample size</i>	n Effect Size 95% CI χ^2 sign.	2 -1.25 -1.53 to -0.97 < 0.0001***	Not required	Not required
Dry weight (g) per plant	<i>Significant, robust but very small sample size</i>	n Effect Size 95% CI χ^2 sign.		2 -0.67 -1.16 to -0.19 0.0061**	2 -0.69 -1.04 to -0.34 < 0.0001***
Height of plants	<i>Not significant, robust but small sample size</i>	n Effect Size 95% CI χ^2 sign.		3 -0.008 -0.89 to 0.87 0.9862 (ns)	3 0.19 -0.69 to 1.07 0.6713 (ns)
Leaves per plant	<i>Not significant, robust but very small sample size</i>	n Effect Size 95% CI χ^2 sign.		2 -0.09 -1.22 to 1.05 0.8814 (ns)	2 -0.06 -1.20 to 1.07 0.9108 (ns)

Table 5: The effectiveness of *Longitarsus jacobaeae* on the reduction of *S. jacobaea*.
n = number of datasets; CI = confidence intervals and χ^2 sign. = chi-squared significance.

Control Measure	Result		Independent/ Non-Independent	Positive Sensitivity Analysis	Negative Sensitivity Analysis
Density of plants	<i>Not significant, robust but small sample size</i>	n Effect Size 95% CI χ^2 sign.	2 -45.67 -139.03 to 47.69 0.3377 (ns)	Not required	Not required
Capitula per plant	<i>Not significant, robust but small sample size</i>	n Effect Size 95% CI χ^2 sign.	2 -31.51 -78.95 to 15.92 0.1929 (ns)	Not required	Not required
Biomass (g) per plant	<i>Not significant, robust but small sample size</i>	n Effect Size 95% CI χ^2 sign.	2 -1.69 -13.20 to 9.82 0.7736 (ns)	Not required	Not required

Table 6: The effectiveness of combining *Tyria jacobaeae* & *Longitarsus jacobaeae*.
n = number of datasets; CI = confidence intervals and χ^2 sign. = chi-squared significance.

Control Measure	Result		Independent/ Non-Independent	Positive Sensitivity Analysis	Negative Sensitivity Analysis
Density of plants	<i>Not significant, robust but small sample size</i>	n Effect Size 95% CI χ^2 sign.	2 -0.086 -1.47 to 1.30 0.9028 (ns)	Not required	Not required

5.4. Time series data

Time Series experiments that studied the control of *S. jacobaea* could not be included within a formal meta-analysis due to the lack of comparator. These datasets are instead presented for each of the natural enemies below in tables 7, 8 and 9, with the addition of the overall reduction of *S. jacobaea* characteristic being calculated for each study.

Table 7: The time series datasets concerning the use of *T. jacobaeae* as a control agent for *S. jacobaea* (ordered in descending percentage reduction).

Study	Time Period (months)	Characteristic	Start	End	% Reduction
Harris, P., Wilkinson, A.T.S. <i>et al.</i> (1976)	36	Stem/m ² (all ages)	0.46	0	100
Cameron, E. (1935)	12	Mature Plants/m ²	26.9	0.0002	99.99
Cameron, E. (1935)	12	Young Plants/m ²	31.5	0.00004	99.99
Harris, P., Wilkinson, A.T.S. <i>et al.</i> (1976)	60	Stem/m ² (all ages)	2.5	0.008	99.68
Nagel, W.P. & Isaacson D.L. (1974)	48	Mature Plants/m ²	3.1	0.4	87.10
Harris, P., Wilkinson, A.T.S. <i>et al.</i> (1976)	48	Rosettes/m ² (all ages)	10	4.2	58
Nagel, W.P. & Isaacson D.L. (1974)	48	Mature Plants/m ²	0.8	0.4	50.00
Nagel, W.P. & Isaacson D.L. (1974)	48	Young Plants/m ²	19.4	11.2	42.27
Nagel, W.P. & Isaacson D.L. (1974)	48	Mature Plants/m ²	4.4	2.8	36.36
Nagel, W.P. & Isaacson D.L. (1974)	48	Mature Plants/m ²	1.8	1.2	33.33
Harris, P., Wilkinson, A.T.S. <i>et al.</i> (1976)	96	Stem/m ² (all ages)	2.7	2	25.92
Nagel, W.P. & Isaacson D.L. (1974)	48	Young Plants/m ²	16.7	13.2	20.96
Nagel, W.P. & Isaacson D.L. (1974)	48	Young Plants/m ²	11.7	9.4	17.09
Nagel, W.P. & Isaacson D.L. (1974)	48	Young Plants/m ²	6.1	5.3	13.11
Harris, P., Wilkinson, A.T.S. <i>et al.</i> (1976)	36	Rosettes/m ² (all ages)	0.03	0.43	increase
Pemberton, R.W. & Turner, C. E. (1990)	72	Plants/m ² (all ages)	53.3	71.1	increase

Table 8: The time series datasets concerning the use of *L. jacobaeae* as a control agent for *S. jacobaea* (all datasets are arranged in descending order)

Study	Time Period (months)	Characteristic	Start	End	% Reduction
M ^c Evoy, P. <i>et al.</i> (1991)	18	Capitula/m ²	160	0	100
M ^c Evoy, P. <i>et al.</i> (1991)	18	Dry mass (g)/m ²	477	0.11	99.9
M ^c Evoy, P. <i>et al.</i> (1991)	18	Plants/m ² (all ages)	308	0.19	99.9
M ^c Evoy, P. <i>et al.</i> (1991)	72	Plants/m ² (all ages)	7.52	0.52	93.1

Table 9: The time series datasets concerning the combined treatment for controlling *S. jacobaea* (all datasets are arranged in descending order)

<i>Tyria jacobaeae</i> & <i>Longitarsus jacobaeae</i> combination					
Study	Time Period (months)	Characteristic	Start	End	% Reduction
Pemberton, R.W. & Turner, C.E. (1990)	216	Plants/m ² (all ages)	15.3	0	100
Pemberton, R.W. & Turner, C.E. (1990)	180	Plants/m ² (all ages)	71	0	100
Hawkes, R.B. & Johnson, G.R. (1976)	72	Plants/m ² (all ages)	15.3	0	100
M ^c Evoy, P.B. (1985)	48	Plants/m ² (all ages)	70	0.6	99.2
Pemberton, R.W. & Turner, C.E. (1990)	180	Plants/m ² (all ages)	11.7	0.18	98.46
M ^c Evoy, P.B. (1985)	36	Dry mass (g)/m ²	718	22	97
Windig, J.J. (1993)	7	Mortality	0	80	80
Windig, J.J. (1993)	7	Mortality	0	60	60

6. DISCUSSION

The purpose of this review was to determine the effectiveness of natural enemies at controlling *S. jacobaea* populations. Using global studies, SMD meta-analyses (Table 4) demonstrates that *T. jacobaeae* significantly reduced certain reproductive characteristics (capitula per plant; seeds per plant / per plot and the viability of seeds) of the *S. jacobaea* plants. The reduction of these characteristics is important in terms of plant resource allocation theory. The results show that *S. jacobaea* plants, in response to herbivory, are possibly allocating more resources to increasing their own survival through their vegetative form (height and rosette size) at the expense of future generations due to their reduced reproductive ability (capitula per plant; seeds per plant / per plot and the viability of seeds). In addition plant resources will also be allocated to the production of pyrrolizidine alkaloids (Pa's) to increase their toxicity and unattractiveness to certain herbivore species attack, (Vrieling & van Wijk, 1994).

There were only 2 datasets for both the *L. jacobaeae* and the *T. jacobaeae* & *L. jacobaeae* combination meta-analyses (Tables 5 & 6). This small sample size increases the risks of generating Type I errors. It is crucial that further high quality primary research is undertaken to allow the effectiveness of these natural enemies to be accurately investigated.

Significantly, the possible reasons for heterogeneity could not be investigated due the small sample sizes within the meta-analyses and, maybe more crucially, the lack of experimental details and site characteristics recorded within the original reports and studies. Future experiments could include the reporting of factors, such as: the soil type / fertility level, which is considered to possibly affect the impact of root feeding insect herbivores such as *L. jacobaeae* (Muller-Scharer, 1991; Steinger & Muller-Scharer 1992). This will allow specific conclusions relating to each factor to be drawn and allow detailed guidance on the effectiveness of each natural enemy under specific site conditions to be available to the practitioner.

The inclusion of time series datasets allows for additional conclusions to be drawn and will assist in the choice of natural enemy in the current absence of higher quality evidence. However, caution is required when using time series as the lower methodological quality allows for confounding factors to creep into the datasets. There was extreme variability for the effectiveness of *T. jacobaeae* (Table 7), with some of the experiments resulting in eradication (no plants left) while others showed an increase in overall *S. jacobaea* densities. Time series datasets for *L. jacobaeae* (Table 8) showed major reductions of *S. jacobaea* densities. However, it was the combination of *T. jacobaeae* & *L. jacobaeae* (Table 9) which showed most consistency in reduction of *S. jacobaea* densities.

The length of follow-up monitoring periods in the included experiments is of concern, with few of those included in the meta-analyses running longer than 12 months. As *S. jacobaea* conforms normally to a biennial nature there might be an initial reduction in the plants densities / characteristic however over a longer follow-up period regeneration of the plant might occur. In future, control projects / experiments should be designed to cover at least a two year time period (hopefully longer).

There are two factors which are crucial in the use of natural enemies in controlling plant populations, especially when using introduced insects. Firstly, the natural enemies have to be able to survive in the environment to which they are introduced. Schmidl (1972) describes experiments on *T. jacobaeae* and its' acclimatisation to new environments. Secondly, the use of natural enemies is unlikely to be successful in countries where there is also a pool of their parasites and predators (e.g. United Kingdom). For a detailed description of the parasites and predators of *T. jacobaeae* see Cameron (1935).

7. REVIEWERS' CONCLUSIONS

7.1. Implications for Conservation/Land Management

The best available evidence suggests that *T. jacobaeae* reduces the reproductive ability of *S. jacobaea* therefore potentially reducing the further spread of the plant. The meta-analysis of *T. jacobaeae* on the densities of *S. jacobaea* showed that the results were not significantly reduced. In addition time series evidence showed vast variability for *T. jacobaeae* effectiveness in controlling *S. jacobaea* densities, with some sites showing total eradication while others had increases.

There were insufficient datasets available to draw any robust conclusions from the meta-analyses for both *L. jacobaeae* and the combination of both natural enemies on *S. jacobaea*. The additional evidence provided by the time series shows that: *L. jacobaeae* caused major reductions of *S. jacobaea* densities and plant characteristics in all datasets. However, it is the use of the combination treatment; of both *T. jacobaeae* & *L. jacobaeae* that shows the greatest potential for the effective control of *S. jacobaea* densities.

The inclusion of time series datasets (those without a comparator), within this systematic review allows for further conclusions to be drawn for all three treatments and will assist the conservation practitioner / policy officer in the choice of which natural enemy to use, until sufficient high quality evidence is available. Caution should be given to these results due to uncertainty of confounding effects and the reduced methodological quality used to obtain the original datasets.

7.2. Implications for Research

Further randomised control trials (RCTs) with multiple replicates and at least two year time periods are required to investigate the effectiveness of all three treatments on *S. jacobaea* densities and plant characteristics. If large sample size and various treatment concentrations (varying the number of natural enemies) are tested, external validity (the generalisations that can be drawn from the observed effects to other populations, settings, or conditions) may be high.

Future trials need vast improvement in the level of details reported on the experimental methodology & results (especially including: number of replicates and the inclusion of a standard deviation measure), along with the site characteristics so that reasons for heterogeneity can be investigated.

8. SOURCES OF SUPPORT

NERC and English Nature grants.

9. ACKNOWLEDGEMENTS

The reviewers would like to thank CEBC team members for their support. Kevin Charman & John Bacon from English Nature and Matthew Oates from The National Trust, for all their support and provision of additional information. We would also like to thank all the practitioners and policy makers who responded to our enquires.

10. REFERENCES

10.1. References to Studies Included in this Systematic Review

- Bornemissza, G.F. (1966) An attempt to control ragwort in Australia with the cinnabar moth (*Callimorpha jacobaeae*)(Arctiidae:Lepidoptera). *Australian Journal of Zoology*, **14**, 201-243.
- Cameron, E. (1935) A Study of the Natural Control of Ragwort (*Senecio Jacobaea* L.). *The Journal of Ecology*, **23**, 265-322.
- Ganeshan, S. (1992) *Direct effect and indirect interactions between flea beetles and cinnabar moths on the ragwort*. (M.Sc and D.I.C.), Imperial College, London, U.K.
- Harris, P. (1974) The impact of the cinnabar moth on ragwort in east and west Canada and its implication for biological control. *Commonw Inst Biol Control*, **8**, 119-123.
- Harris, P., Thompson, L.S., Wilkinson, A.T.S., & Neary, M.E. (1976) Reproductive biology of tansy ragwort, climate and biological control by the cinnabar moth in Canada. In IV International symposium on biological control of weeds, pp. 163-173, Gainesville, Florida, USA.
- Harris, P., Wilkinson, A.T.S., Thompson, L.S., & Neary, M.E. (1976) Interaction between the cinnabar moth, *Tyria jacobaeae* and ragwort, *Senecio jacobaea* in Canada. In IV International symposium on biological control of weeds, pp. 174-180, Gainesville, Florida, USA.
- Hawkes, R.B. & Johnson, G.R. (1976) *Longitarsus jacobaeae* Aids Moth in the Biological Control of Tansy Ragwort. In Proceedings of the 4th International Symposium on Biological Control of Weeds, pp. 193-196.
- James, R.R., McEvoy, P.B., & Cox, C.S. (1992) Combining the Cinnabar Moth (*Tyria-Jacobaeae*) and the Ragwort Flea Beetle (*Longitarsus-Jacobaeae*) for Control of Ragwort (*Senecio-Jacobaea*) - an Experimental-Analysis. *Journal of Applied Ecology*, **29**, 589-596.
- McEvoy, P.B. (1985) Depression in ragwort (*Senecio jacobaea*) abundance following introduction of *Tyria jacobaea* and *Longitarsus jacobaeae* on the central coast of Oregon. *Proceedings of the VI International Symposium on Biological Control of Weeds / edited by Ernest S. Delfosse. Ottawa : Agriculture Canada*, 57-64.
- McEvoy, P., Cox, C., & Coombs, E. (1991) Successful Biological Control of Ragwort, *Senecio Jacobaea*, by Introduced Insects in Oregon. *Ecological Applications*, **1**, 430-442.
- Nagel, W.P. & Isaacson, D.L. (1974) *Tyria jacobaea* and tansy ragwort in western Oregon. *Journal of economic entomology*, **67**, 494-496.
- Pemberton, R.W. & Turner, C.E. (1990) Biological-Control of *Senecio-Jacobaea* in Northern California, an Enduring Success. *Entomophaga*, **35**, 71-77.
- Schmidl, L. (1972) Studies on the control of ragwort, *Senecio jacobaea* L., with the cinnabar moth, *Callimorpha jacobaeae* (L.) (Arctiidae: Lepidoptera), in Victoria. *Weed research*, **12**, 46-57.

Windig, J.J. (1993) Intensity of Longitarsus-Jacobaeae Herbivory and Mortality of Senecio-Jacobaea. *Journal of Applied Ecology*, **30**, 179-186.

10.2. References to Studies Excluded from this Systematic Review

All references viewed at both the title and abstract or at the full text filtering stages are available on request as either an Endnote Library or Excel file.

10.3. Additional References

Egger, M., Smith, G.D., & Altman, D.G., eds. (2003) *Systematic Reviews in Health Care: Meta-analysis in context*, 2nd edn. BMJ Publishing Group, London.

Frick, K.E. (1971) *Longitarsus jacobaeae*, a flea beetle for the biological control of tansy ragwort II. *Ann. Entomol. Soc.*, **64**, 834.

Frick, K.E. & Johnson, G.R. (1973) *Longitarsus jacobaeae*, a flea beetle for the biological control of tansy ragwort 4: Life history and adult aestivation of an Italian biotype. *Ann. Entomol. Soc.*, **66**, 358-367.

Muller-Scharer, H. (1991) The impact of root herbivory as a function of plant density and competition: survival, growth and fecundity of *Centaurea maculosa* in field plots. *Journal of Applied Ecology*, **28**, 759-776

Steinger, T. & Muller-Scharer, H. (1992) Physiological and growth responses of *Centaurea maculosa* (Asteraceae) to root herbivory under varying levels of interspecific plant competition and soil nutrient availability. *Oecologia*, **91**, 141-149.

Sutton, A.J., Abrams, K.R., Jones, D.R., Sheldon, T.A., & Song, F. (2000) *Methods for Meta-Analysis in Medical Research* John Wiley & Sons, Chichester.

Vrieling K., van Wijk C.A.M. (1994) Cost assessment of the production of pyrrolizidine alkaloids in ragwort (*Senecio jacobaea* L.). *Oecologia*, **97**, 541-546.

Appendix 1. Quality Assessment Instrument to provide an estimate of bias surrounding extracted data.

Bias	Generic data quality features	Specific data quality features	Quality element	Quality score
Selection and Performance bias	Study Design	NA	Randomized controlled Trial	80
			Quasi-RCT (a trail applying a pseudo random allocation mechanism, e.g. date of planting)	70
			Controlled Trial	60
			Historical CT (data for the control arm comes from archives or is calculated, therefore not from current experimental observation)	50
			Site comparison	40
			Time Series	30
			Interrupted time series	20
			Questionnaire	10
			Expert Opinion	10
	Baseline comparison (heterogeneity between treatment and control arms with respect to defined confounding factors before treatment)	Factors: Size of experimental area	Treatment and control arms homogenous	1
			Treatment and control arms not comparable with respect to confounding factors OR insufficient information	0
		Habitat type,	Treatment and control arms homogenous	1
			Treatment and control arms not comparable with respect to confounding factors OR insufficient information	0
		Location/Geographical Area	Treatment and control arms homogenous	1
			Treatment and control arms not comparable with respect to confounding factors OR insufficient information	0
Altitude.	Treatment and control arms homogenous	1		

	Baseline comparison (cont.)		Treatment and control arms not comparable with respect to confounding factors OR insufficient information	0		
		Plant age at time of treatment	Treatment and control arms homogenous	1		
			Treatment and control arms not comparable with respect to confounding factors OR insufficient information	0		
		Soil Type	Treatment and control arms homogenous	1		
			Treatment and control arms not comparable with respect to confounding factors OR insufficient information	0		
		Intra treatment variation	Factors: Plant age at time of treatment	No heterogeneity within treatment and control arms	1	
	Replicates within treatment and control arms not comparable			0		
	Habitat type,		No heterogeneity within treatment and control arms	1		
			Replicates within treatment and control arms not comparable	0		
	Location,		No heterogeneity within treatment and control arms	1		
			Replicates within treatment and control arms not comparable	0		
	Altitude.		No heterogeneity within treatment and control arms	1		
			Replicates within treatment and control arms not comparable	0		
	Measurement of intervention and Co-interventions		Biological control agent	Factor equal in treatment and control	1	
				Factor not equal or unreported	0	
	Assessment bias		Measurement of outcome	Replication, parameter of abundance (accuracy)	Well replicated objective parameter of abundance used (>5 replications)	4
					Replicated objective parameter of abundance used (>2 replications)	2
		Unreplicated however parameter measured sensible			1	
Unreplicated observations or subjective parameter of abundance used		0				

Appendix 2. Study Characteristics Tables for each study accepted at full text.

Study 1	Bornemissza, G. F. (1966)												
Methods	A control trial (CT) on an abandoned dairy farm, comparing defoliated plants (by cinnabar moths) against control (moth excluded).												
Population	Size of experimental area:						Unknown size of area – however a total of 200 plants (100 in each arm), all contained within sleeving were used in the experiment.						
	Habitat:						Abandoned dairy farm pasture fields – reverting to weed & scrub.						
	Location:						nr. Gunyah, Victoria, Australia.						
	Altitude:						approx. 365m a.s.l.						
	Plant age at time of treatment:						Unknown, however all of similar age and size within & between experimental arms.						
	Soil type:						Heavy-textured soils.						
Weather	Data from 3km from the study site (Olsen Bridge, Victoria, Australia.). The region has an average annual rainfall of 1371mm, well distributed over the year but 1016mm falling within the months of May-November. Drought conditions are frequent in summer, while winter is foggy and cold with mild morning ground-frosts. The values are the means for the decade 1952-61.												
	Observation	J	F	M	A	M	J	J	A	S	O	N	D
	Max. air temp (°C)	23	22	21	18	14	12	11	12	14	16	18	20
	Rainfall (mm)	61	67	66	87	160	181	151	145	139	141	139	109
	Hot days (+27°C)	11	7	6	3	-	-	-	-	-	1	4	5
	Rainy days	9	11	12	14	21	21	21	22	18	19	18	14
Intervention & Comparator	Each arm of the experiment has 100 ragwort plants of similar age and size (200 plants in total). Organdie sleeves are used to either enclose 25 <i>Tyria</i> (cinnabar moth caterpillars) on ragwort plants (Treatment group) or to exclude them (Control group). The dry weight (g), capitula per plant, seed per capitula, and seed per plant were recorded. Values are means of 100 plants = 33 plants after defoliation, 66 plants after regeneration. Control values marked with * are from the beginning of January. The values for Effective Reduction are taken from the paper. Measurements were made simultaneously – after defoliation = end of January; after regeneration = mid-April, after the plants have had time to regenerate after <i>Tyria</i> attack.												
Outcomes							Treatment		Control		Effective Reduction		
	Dry Weight (g)		After defoliation				16		33		33%		
			After regeneration				17		34				
	Capitula per Plant		After defoliation				0		560		83%		
			After regeneration				87		565				
	Seed per Capitula		After defoliation				0		75*		43%		
			After regeneration				43						
	Seed per Plant		After defoliation				0		42280*		88%		
			After regeneration				3741						
	Viability of Seeds (%) after a 4 month follow-up								Treatment =		39%		
								Control =		52%			

Study design	Control Trial	60pts
Baseline comparison	All factors known (Size, Habitat, Location, Altitude, Plant Age and Soil)	6pts
Intra treatment variation	All factors again know (Age, Habitat, Location, Altitude)	4pts
Measurement of intervention and co-interventions	All factors know (bio-control agent & number, no other interventions)	2pts
Replication & parameter of abundance	Replication even for each experimental arm, with outcome parameter measured equally and simultaneously between arms.	4pts
Data Quality Score Total		76pts
Other notes	<p>Data extracted from Table 2 of the original paper.</p> <p>Plants that were attacked / defoliated usually produced vigour secondary growth and a belated crop of seeds.</p> <p>Larvae feed more voraciously during their last two instars. This occurred at the study site at the height of the ragwort flowering season. Thus, larval attack will cease and defoliation will normally have been effected, before all the small buds have developed into flowering capitula and while most plants still possess substantial reserves.</p>	

Study 2	Cameron, E. (1935)								
Methods	A historical (estimated) control trial (HCT) using estimated seed yields based on a calculation from previous experiments to calculate the control arm of the experiment. Also a timeseries of the effectiveness of Tyria on decreasing ragwort plant densities.								
Population	Size of experimental area:			Varies with each area – see below					
	Habitat:			Impoverish pastures					
	Location:			Varies – see below					
	Altitude:			Unknown					
	Plant age at time of treatment:			Given by plant height – see below					
	Soil type:			Unknown					
Weather	No weather data was presented / recorded for any of the sites within this study.								
Intervention & Comparator	Datasets A & B A HCT based on a calculation of estimated seed yield for the control arm of the experiment. Each of the experimental areas varies in size, number of plants and number of Tyria attacking the ragwort. All plants = common ragwort (<i>S. jacobaea</i>). n is based on number of plants								
Outcomes	Area ID	Plot Size (m²)	Number of Plants	Av. Height of plants (cm)	Number of Tyria	Treatment (av. Seed yield per plant)	Control (est. Seed yield per plant)	Length of follow-up (months)	
	Wentworth (site A)	1.5	14	61	15	974	20000	2	
	Stoke Poges (B)	114	14	70	15	0	23689	4	
Intervention & Comparator	Dataset C A timeseries showing the effectiveness of Tyria at controlling ragwort plant densities – when secondary growth, owing to an unfavourable session, was eliminated. Record from fields at Fawley Court Hill, Henley, U.K. 1932. Timeseries length = 12 months. Area of the field = 4.86 hectares, values below have been recalculated from per acre. Average of 12 Tyria on each plant observed.								
Outcomes		Date	Mature Plants		Young Plants				
			Per hectare	Total field	Per hectare	Total field			
	Start	June 1931	269278	1306800	315364	1529520			
	End	June 1932	2.47	12	0.41	2			
	Change		99.99%	99.99%	99.99%	99.99%			
Study design	2 historical control trial (HCT) (datasets A-B)							50pts	
	1 timeseries (dataset C)							30pts	
Baseline comparison	Unknown factors include – altitude of the site & soil type.							All datasets = 4pts	
Intra treatment variation	Again unknown factors included – altitude.							All datasets = 3pts	
Measurement of intervention and co-interventions	Tyria used as Bio-control agent – the number per plant was recorded for all sites and no other co-interventions were being undertaken.							All datasets = 2pts	
Replication & parameter of abundance	All HCT experiments were well replicated and even between each arm of the experiment.							4pts	
	No replication seemed to be undertaken							0pts	
Data Quality Score Total	Datasets A-B				Dataset C				
	63pts				39pts				
Other notes	Paper draws the following conclusions: 1. Providing that the attack is general & that no secondary growth follows, either in the shape of a new crop of flowers, or as new growth from base, to carry the ragwort over into another year, Tyria can be a very effective control agent. 2. When poor plants, growing on very inferior soil, are heavily and uniformly attacked by Tyria the ragwort infestation should be wiped out. 3. Once an infestation is under control, certain precautions have to be taken in order that the area may be kept free of ragwort. These take the form of (a) stimulation of grasses in the area, (b) avoidance of overgrazing.								

Study 3	Ganesham (1992)				
Methods	A Control trial investigating the effect of cinnabar moth herbivory on density of plants, number of capitula produced and mean height of flowering plants.				
Population	Size of experimental area:	Plots of 10m ² (0.001 hectares)			
	Habitat:	Field			
	Location:	Silwood Park, Berkshire, U.K. (Grid Ref: 41/944 691)			
	Altitude:	Unreported – sloped field			
	Plant age at time of treatment:	Varies naturally random – no set age			
	Soil type:	Unreported			
Weather	No weather data was presented.				
Intervention & Comparator	<p>Control Trial consisting of 3 factors with two levels each as follows:</p> <ol style="list-style-type: none"> 1. Cinnabar moth +/- 2. Rabbits +/- 3. Soil Insecticide +/- <p>The cinnabar moths were hand picked from plots to remain free from damage and transferred to those plots where cinnabar moths were meant to be present. Rabbit grazing was prevented by fencing. Soil insecticide, Dursban (chlorpyrifos), was applied from autumn 1991 every 6-8 weeks. 4 replicates were taken for the experiments. The density of plants, number of capitula produced and mean height of flowering plants were recorded for each plot. (The impact of insecticides and rabbits could not be investigated as the other contributing data to the meta-analysis did not have sufficient detail to allow investigation).</p>				
Outcomes (density of ragwort plants)		Dataset A		Dataset B	
	Cinnabar	Insecticide		No soil insecticide	
		+ Rabbit	- Rabbit	+Rabbit	- Rabbit
	With (treatment)	3.25	0.25	5.00	0.25
	Without (control)	6.25	0.25	2.25	1.00
Outcomes (mean number of capitula per plant)		Dataset A		Dataset B	
	Cinnabar	Insecticide		No soil insecticide	
		+ Rabbit	- Rabbit	+Rabbit	- Rabbit
	With (treatment)	9.364	2.00	37.216	0
	Without (control)	14.130	0	16.667	42.577
Outcomes (mean height (cm) of flowering plants)		Dataset A		Dataset B	
	Cinnabar	Insecticide		No soil insecticide	
		+ Rabbit	- Rabbit	+Rabbit	- Rabbit
	With (treatment)	43.82	77.00	48.08	66.00
	Without (control)	46.22	72.00	32.33	60.19
Study design	A control trial				60pts
Baseline comparison	Altitude & soil type are unknown				4pts
Intra treatment variation	Altitude varies as the field is on a slope & soil type/characteristics are unknown				3pts
Measurement of intervention and co-interventions	Co-interventions are known (rabbits & soil insecticide) and are factored into account with experimental design. However unknown number of Tyria per plot.				1pts
Replication & parameter of abundance	4 replicates – measures are all sensible				2pts
Data Quality Score Total	For both datasets A & B		70pts		
Other notes					

Study 4	Harris, P. (1974)	
Methods	Site Comparison of cinnabar moth herbivory on ragwort at Nova Scotia, Canada.	
Population	Size of experimental area:	Unreported
	Habitat:	Pasture fields and roadsides
	Location:	Nova Scotia & Prince Edward Island, Canada.
	Altitude:	Unreported
	Plant age at time of treatment:	Immature plants
	Soil type:	Unreported
Weather	No weather data was presented within the study.	
Intervention & Comparator	A site comparison study, with records of the ragwort plants density m² for the year 1972 at Nova Scotia, Canada. The comparison site was the nearest permanent pasture which was unaffected by the cinnabar moth (<i>Tyria</i>) defoliation. The number taken for replication = 1.	
Outcomes (Control Trial)		Immature plants m²
	Treatment	0.03
	Control	62.2
Study design	Site Comparison	40pts
Baseline comparison	Size of the experimental area, Altitude of sites and soil type are all unreported. The plant ages are all immature.	3pts
Intra treatment variation		
Measurement of intervention and co-interventions	No co-interventions reported. However, an unknown number of <i>Tyria</i> per plot/plant.	1pt
Replication & parameter of abundance	Number of replicates unknown therefore only 1 used. Sensible measure of density (m ²)	1pt
Data Quality Score Total		45pts
Other notes	Ragwort recorded as being perennial within Canada – most perennial plants can tolerate a single defoliation with few ill effects unless it is timed to force the plant into a period of stress such as winter or drought while it is in a physiologically unstable state. Therefore, it is worth-while making an effort to determine this period before introducing an agent to control a perennial weed. However, if the climate is equitable throughout the year, the only strategy that will succeed is to attack the weed throughout the growing season. This can be achieved by a multivoltine agent or a series of univoltine agents.	

Study 5	Harris, P., Thompson, L. S., Wilkinson, A. T. S. & Neary, M. E. (1976) Known as: Harris, P., Thompson, L. S. <i>et.al.</i> (1976)			
Methods	A site comparison study with 4 sites defoliated by cinnabar moths and 1 site left as an untreated control			
Population	Size of experimental area:	Each plot was 1m ²		
	Habitat:	Pasture fields		
	Location:	Canada (Nanaimo, British Columbia & Durham, Nova Scotia).		
	Altitude:	Unreported		
	Plant age at time of treatment:	Both rosettes & flowering plants		
	Soil type:	Unreported		
Weather	No weather was reported within the study.			
Intervention & Comparator	<u>Site comparison data</u> Both flowering plants & rosettes on two 1m ² plots at Nanaimo, B.C. were tagged & the number of leaves on each rosette were recorded. The plants were then defoliated by adding several hundred field-collected cinnabar larvae. One year later the survival of each plant was noted and the numbers of leaves on the rosettes were counted. A similar procedure was followed at Durham, N.S. except that one plot was left un-defoliated as a control site. The follow-up time was 12 months with the experiment ending in 1973. An average of 3 cinnabar moth larvae to plant was calculated. Again the number of replication = 1.			
Outcomes	Dataset	Number of rosettes (density m²)	Number rosette leaves/plot	Average leaves/plant
	A	183	612	3.255
	B	71	244	3.437
	C	59	336	5.695
	D	3	0	0
	Control	72	541	7.514
Study design	Study comparison study			40pts
Baseline comparison	Soil type/characteristics and the altitudes of each site are unknown. All other factors reported and similar.			4pts
Intra treatment variation	Only details of altitude of each of the sites unreported.			3pts
Measurement of intervention and co-interventions	The average number of Tyria per plant could be approximately calculated from the text. No co-interventions reported on sites during the experiment.			2pts
Replication & parameter of abundance	No replication of results only one plot per field recorded. Sensible parameter of assessment was used.			1pt
Data Quality Score Total	For datasets A-D		50pts	
Other notes				

Study 6	Harris, P., Wilkinson, A. T. S., Thompson, L. S. & Neary, M. E. (1976) Known as: Harris, P., Wilkinson, A. T. S. <i>et.al.</i> (1976)				
Methods	Three different timeseries measuring ragwort/m ² against cinnabar moth larvae/m ² for the east and west coasts of Canada.				
Population	Size of experimental area:	Varies – see below.			
	Habitat:	Farmland pastures and abandoned fields.			
	Location:	Three sites in Canada as follows: a) Durham, Nova Scotia b) Selkirk, Prince Edward Island c) Nanaimo, British Columbia			
	Altitude:	Unreported			
	Plant age at time of treatment:	All ages – as measuring stems/m ²			
	Soil type:	Varies with site: a) Unreported b) Sandy loam (poor drainage) c) Gravelly sandy loam			
Weather	No weather was reported within the study.				
Intervention & Comparator	<u>Dataset A – Durham, Nova Scotia</u> Release field = 3,500m ² with a dense stand of ragwort since 1955, caused by over grazing and uneven ground. Recording was undertaken during mid-July, allowing the majority of eggs to hatch and the ragwort to be bloom. Rosettes m ² were also recorded for some of the timeseries.				
Outcomes	Year	Stems/m²	Rosettes/m²	Larvae/m²	Larvae/stem
	1969	2.5		8.0	3.2
	1970	0.25		1.4	18.0
	1971	0.06		0.6	10.0
	1972	0.001	0.03	0.09	90.0
	1973	0.008	0.06	0.004	0.5
	1974	0.008	0.43	0.008	1.0
Intervention & Comparator	<u>Dataset B – Selkirk, Prince Edward Island</u> Release field = 26,200m ² (2.62ha) of abandoned farmland with a continuous sward of grass & mouse-eared hawkweed (<i>Hieracium pilosella</i>) and was not cut or grazed. The flowering stems shorter and the density of the ragwort was considerably less than at Durham; however heavy stands of infestation were present. Sampling completed during mid-July, the week after that for Durham, as emergence of larvae was later than that in Nova Scotia. Ragwort densities calculated at paced intervals across the field.				
Outcomes	Year	Stem/m²	Larvae/m²	Larvae/stem	
	1971	0.46	1.2	2.6	
	1972	0.02	0.1	5.0	
	1973	0.0	0.0	0.0	
Intervention & Comparator	<u>Dataset C – Nanaimo, British Columbia</u> Release field = 75,000m ² (7.5ha) with an easterly exposure. Much of the field was subject to severe summer drought, which usually defoliated ragwort rosettes. Field was surrounded by Douglas fir (<i>Pseudotsuga menziesii</i>), hemlock (<i>Tsuga heterophylla</i>) and western red cedar (<i>Thuja plicata</i>). Sampling occurred over a period of June to July for the moth with the ragwort stems being counted over the summer. Rosettes m ² were also recorded for some of the timeseries.				
Outcomes	Year	Stems/m²	Rosettes/m²	Larvae/m²	Larvae/stem
	1968	2.7		10.4	3.8
	1969	2.0		8.0	4.1
	1970	2.5		6.5	5.2
	1971	1.4		2.5	5.3
	1972	2.1	10.0	1.2	2.3
	1973	0.5	14.9	0.5	1.0
	1974	0.4	5.0	0.2	0.5
1975	2.0	4.2	0.1	0.05	

Study design	All 3 are datasets are timeseries		30pts
Baseline comparison	Altitude and soil type is unreported All factors similar except altitude which is unreported.		(Dataset A) 4pts (Datasets B-C) 5pts
Intra treatment variation	All datasets have missing altitude information; all other factors are known and similar.		3pts
Measurement of intervention and co-interventions	Co-interventions such as grazing is still on-going.		2pts
Replication & parameter of abundance	Unknown level of replication – i.e. number of samples taken to calculate value m ² . A sensible parameter of assessment was used.		1pt
Data Quality Score Total	Dataset A		40
	Dataset B – C		41
Other notes			

Study 7	Hawkes, R. B. & Johnson, G. R. (1976)				
Methods	A timeseries and two control trials using the ragwort flea-beetle (<i>Longitarsus jacobaeae</i>) as a singular or combined control agent with cinnabar moth larvae across three sites in Fort Bragg, California, U.S.A.				
Population	Size of experimental area:	Varies with each site – see below.			
	Habitat:	River bottom pastures			
	Location:	Fort Bragg, California, U.S.A.			
	Altitude:	Unreported			
	Plant age at time of treatment:	All ages			
	Soil type:	Unreported			
Weather	No weather was reported within the study.				
Intervention & Comparator	Dataset A – Timeseries – Site 1 The original release site for <i>L. jacobaeae</i> in the Fort Bragg region. It is ca. 5 ha of river bottom pasture with heavy grass vegetation which originally contained ragwort. Grazed by sheep, sometimes feeding on ragwort in small quantities. Plant density was recorded with a 1m ² quadrat, every five paces along a transect. Based on a 72 month (6 year) timeperiod.				
Outcomes			Ragwort density/m²		
	Start (1969)		15.3		
	End (1975)		0.0		
Intervention & Comparator	Dataset B & C – Control Trial – Site 2 & 3 Site 2 – has not been grazed for years, with a dense stand of velvetgrass (<i>Holcus lanatus</i>) & several spp of annual & perennial forbs. In October 1972 ca. 2000 flea beetles were released on the site. A control site 500m to the east was established where just cinnabar moths were naturally present (this distance was required due to the dispersal of <i>L. jacobaeae</i>). Site 3 – ca. 5 ha, it is closely grazed by cattle, which ignore feeding on the ragwort. Ca. 5000 <i>L. jacobaeae</i> were released in October 1972. A control plot was established 200m to the west of the treatment site. Data was extracted for a follow-up period of 2 years prior to <i>L. jacobaeae</i> dispersing to the control site. The average <i>L. jacobaeae</i> was 31.7 larvae/rosette. Data was extracted and converted to reduction of ragwort /m ² . Replication, n = 1 for both datasets.				
Outcomes	Densities (plant/m²)				
		Site 2 (dataset B)		Site 3 (dataset C)	
		Treatment	Control	Treatment	Control
	Start	71.1	36.1	18.3	7.7
	End	6.9	14.5	7.4	9.0
	Overall reduction	64.2	21.6	10.9	-1.3 (increased density)
Study design	Timeseries Control Trial		(dataset A) 30pts (datasets B-C) 60pts		
Baseline comparison	Altitude and soil type is unreported		(all datasets) 4pts		
Intra treatment variation	Only altitude is unreported, all other factors are similar		(all datasets) 3pts		
Measurement of intervention and co-interventions	Co-interventions such as grazing are reported and is still on-going at sites 1 & 3.		(all datasets) 2pts		
Replication & parameter of abundance	Unknown level of replication to obtain data – parameter of abundance is suitable.		(all datasets) 1pt		
Data Quality Score Total	Dataset A		40		
	Datasets B – C		70		
Other notes					

Study 8	James, R. R., McEvoy, P. B. & Cox, C. S. (1992) Known as: James, R. R., et al. (1992)				
Methods	An exclusion experiment (RCT) was conducted using cages & experimental ragwort populations to determine which of the following was most effective in depressing plant populations: <i>Tyria jacobaeae</i> , <i>Longitarsus jacobaeae</i> , or a combination of both.				
Population	Size of experimental area:	0.9 ha divided into four blocks each with 15 plots of 0.25m ² .			
	Habitat:	Meadow			
	Location:	Cascade Head Scenic Research Area, central coast Oregon. U.S.A.			
	Altitude:	Unreported			
	Plant age at time of treatment:	All ages – from transplanted plants & seed			
	Soil type:	Unreported			
Weather	No weather was reported within the study.				
Intervention & Comparator	<p>Meadow was divided into four blocks. 15 plots of 0.25m² were placed within a randomly located 3*4m area in each of the blocks. Natural vegetation was removed from the meadow other than ragwort and ragwort transplants were thinned to recreate the situation of when the biocontrol agents were introduced to the area. Insects were excluded from the area via cages for 1 year, while the plants became established (flowering plants and seedlings). Experimental plots were covered with 61*61*61cm frames constructed of 2.5cm diameter plastic (PVC) tubes covered with bags of 'Leno weave' nylon mesh screens. Each plot was assigned to one of five treatments:</p> <ol style="list-style-type: none"> 1. Neither insect (control) 2. Moth only – cages which excluded flea-beetle but opened June – July for cinnabar moth; flowering plant defoliated. 3. Beetle only – cages open to flea-beetle but closed June – July to exclude moth. 4. Both insects – cages continuously open. 5. Open control – to see the side effects of the cages. <p>Each of the treatments were replicated three times, within each block to allow for two destructive samples and a census plot, results are a mean of 4 plots. The experiment was randomised block design. Some flea-beetles did manage to enter cages however less than 1% of that in the beetle only treatment and 1.2% of total beetles in the both insect treatment. For further details please see original study.</p>				
Outcomes		Control	Tyria only (A)	Longitarsus (B)	Both (C)
	Ragwort Density	53.25	50.37	43.1	
	Total Biomass (g/plot)	165.3	149.7	207.5	167.0
	Beetle larvae per plot	4.9		322.8	251.1
	Leaves per stem	37.9	37.0	31.5	8.8
	Capitula per stem	240.6	53.9	144.0	2.4
	Achenes per capitula	65.5	55.0	66.5	53.5
	% Viable Achenes	34.9	25.7	54.3	0.0
Study design	Randomised control trial (RCT)				80pts
Baseline comparison	Altitude and soil type is unreported				4pts
Intra treatment variation	Only altitude is unreported, all other factors are similar				3pts

Measurement of intervention and co-interventions	No co-intervention on-site, excellent recording of biocontrol methods		2pts
Replication & parameter	Replicated experiment with suitable parameters of abundance reported		
Data Quality Score Total	For Dataset A-C	91pts	
Other notes	<p>Taken from the Discussion:</p> <p>“...support the hypothesis that two insects together, feeding on different plant parts & at different times of the year, can have a greater impact on host-plants than either insect acting alone.”</p> <p>“Alone, beetles decreased vegetative plan biomass & density and the cinnabar moth treatment reduced fecundity.”</p> <p>“The combination of both herbivore treatments reduced achene production & viability to the extent that fecundity was negligible.”</p> <p>“The high mortality of young plants (80-99%) caused by beetle activity undermined the pyramidal structure of the plant population, leaving fewer individuals to be recruited into the reproductive stage.”</p>		

Study 9	McEvoy, P. B. (1985)	
Methods	A timeseries reporting the number releases of cinnabar moth and flea beetle comparing it to the annual estimates of standing crop (dry g/m ²).	
Population	Size of experimental area:	0.9 ha
	Habitat:	Abandoned diary farm pasture
	Location:	Cascade Head Scenic Research Area, central coast Oregon. U.S.A.
	Altitude:	Unreported
	Plant age at time of treatment:	All ages
	Soil type:	Unreported
Weather	No weather was reported within the study.	
Intervention & Comparator	Timeseries A Total release of biocontrol agents at the site included 2000 cinnabar moths in 1978 and 230 flea-beetles in 1979 and an additional 485 in 1980. Biocontrol agent's populations remained low until 1982/3 when a population explosion occurred. Four years worth of data was recorded.	
Outcomes		Ragwort density/m²
	Start (1980)	71
	End (1984)	0.6
	Total decline (over 48 months)	70.4 (99.2% reduction)
Intervention & Comparator	Timeseries B The standing crop (dry g/m ²) is recorded using quadrats randomly placed over the site. This is based on 3 years worth of records	
Outcomes		Standing ragwort crop (dry g/m²)
	Start (1981)	718
	End (1984)	22
	Total decline (over 36 months)	696 (97% reduction)
Study design	Timeseries	30pts
Baseline comparison	Altitude and soil type is unreported	4pts
Intra treatment variation	Only altitude is unreported, all other factors are similar	3pts
Measurement of intervention and co-interventions	No co-intervention on-site, previous activities on site (dairy farming). Biocontrol methods and numbers introduced are reported	2pts
Replication & parameter of abundance	No replication as timeseries	0pts
Data Quality Score Total	Datasets A-B	39pts
Other notes	While the ragwort crop declined – the standing crop of other species increased by 10 fold (707 dry g/m ²). This lead to a no net change in mean standing crop of the plant communities.	

Study 10	McEvoy, P., Cox, C. & Coombs, E. (1991) Known as: McEvoy, P., et al (1991)			
Methods	A number of Randomised Control Trials and Timeseries studying the response of ragwort to <i>Longitarsus jacobaeae</i> defoliation.			
Population	Size of experimental area:	0.9 ha with plots 0.25m ²		
	Habitat:	Abandoned diary farm pasture		
	Location:	Cascade Head Scenic Research Area, central coast Oregon. U.S.A.		
	Altitude:	Only for dataset C – see appendix of original study		
	Plant age at time of treatment:	All ages		
	Soil type:	Unreported		
Weather	Average precipitation mm/yr given for dataset C in appendix of the original study.			
Intervention & Comparator	<p><u>Dataset A – Perturbation Experiment</u></p> <p>The objective of this experiment was to “create high-density ragwort populations & then to compare ragwort density, biomass & reproduction (capitulum) after the plants were exposed to & protected from introduced biological control trial.”</p> <p>A single-factor randomised-block design with two treatments (treat & control), four blocks & 1 replication per block. Beetles were removed from the control (caged) plots over a period of a year before experiment started and insecticide was sprayed during the experiment so the plots remained clear. Most of the values below have been scaled up to value/m² and read from figure 3 in the original paper. The beetle density was 906/per plot or 3624/per m².</p>			
Outcomes	End Levels – August 88			
		Treatment	Control	
	Density/m²	0.76	1000	
	Dry mass (g)/m²	0.44	80	
	Capitula/m²	0	600	
Intervention & Comparator	<p><u>Dataset B – Perturbation Experiment (timeseries)</u></p> <p>As above – a timeseries of 18 months.</p>			
Outcomes		Start	End	% change
	Density/m²	308	0.76	-99.9
	Dry mass (g)/m²	477	0.44	-99.9
	Capitula/m²	160	0	-100
Intervention & Comparator	<p><u>Dataset C – site comparison</u></p> <p>Follow-up of 6 years (72 months) measuring the ragwort densities/m² following the release of <i>Longitarsus jacobaeae</i>. Data is based on 14 sites across Oregon</p>			
Outcomes		Start	End	% change
	Density/m²	7.52	0.52	-93.1
Study design	Randomised Control Trial		(dataset A) 80pts	
	Timeseries		(datasets B-C) 30pts	
Baseline comparison	Altitude and soil type is unreported		(datasets A-B) 4pts	
	Only soil type unreported – Altitude is given for each site		(dataset C) 5pts	
Intra treatment variation	All other factors are similar – expect altitude		(all datasets) 3pts	
Measurement of intervention and co-interventions	All factors known for the site		(datasets A-B) 2pts	
	Co-interventions are unknown for all the sites		(dataset C) 1pt	
Replication & parameter of abundance	Replicated dataset with suitable parameter measured		(dataset A) 2pts	
	No replication, however suitable parameter is measured		(dataset B-C) 1pt	
Data Quality Score Total	Dataset A		91 pts	
	Dataset B		40 pts	
	Dataset C		40 pts	

Study 11	Nagel, W. P. & Isaacson, D. L. (1974)				
Methods	A timeseries of four sites studying the impact that larvae of <i>Tyria jacobaeae</i> have on common ragwort.				
Population	Size of experimental area:	Unreported – However results listed as m ²			
	Habitat:	Varies – see below			
	Location:	Myrtle Point, Oregon, U.S.A.			
	Altitude:	Varies – see below			
	Plant age at time of treatment:	Varies – studies both young & flowering plants			
Soil type:	Unreported				
Weather	A mild climate with wet winters and dry summers.				
Intervention & Comparator	<p>The four study sites were located south of Myrtle Point, Oregon, U.S.A.</p> <p>Site 1 – improved pasture; 700ft elevation; plant cover of grass, forbs & ragwort.</p> <p>Site 2 – prairie pasture; 650ft elevation; plant cover of grass, forbs & ragwort.</p> <p>Site 3 – cleared pasture; 160ft elevation; grasses, poison oak patches & ragwort.</p> <p>Site 4 – semi-cleared pasture; 500ft elevation; grasses, forbs, ragwort, Douglas-fir, white fir, red alder & big-leaf maple.</p> <p>Sampling was undertaken in June/July of each year.</p>				
Outcomes Site 1		1970	1971	1972	1973
	1 st year plants/m ²	19.4±5.4	20.9±7.6	27.2±6.0	11.2±6.0
	Flowering plants/m ²	4.4±0.9	4.6±1.1	3.6±0.8	2.8±1.3
	Biomass (dry g/m ²)	20.2±3.4	23.2±4.6	16.4±2.6	9.7±4.1
Tyria density/m ²	0.7±0.3	5.4±3.0	1.5±0.9	16.5±7.1	
Outcomes Site 2		1970	1971	1972	1973
	1 st year plants/m ²	11.7±1.5	11.1±1.5	9.3±1.8	9.4±1.9
	Flowering plants/m ²	3.1±0.9	0.9±0.3	0.9±0.3	0.4±0.2
	Biomass (dry g/m ²)	15.8±3.8	9.4±2.1	8.8±2.8	3.6±0.7
Tyria density/m ²	1.3±0.7	2.8±1.0	1.6±1.3	1.5±0.6	
Outcomes Site 3		1970	1971	1972	1973
	1 st year plants/m ²	6.1±2.1	8.2±1.6	9.4±3.0	5.3±2.6
	Flowering plants/m ²	1.8±0.5	1.9±0.5	2.2±0.9	1.2±0.5
	Biomass (dry g/m ²)	8.4±2.8	10.3±2.2	7.3±2.5	2.7±1.3
Tyria density/m ²	9.8±6.0	8.4±2.4	6.2±3.2	4.2±1.3	
Outcomes Site 4		1970	1971	1972	1973
	1 st year plants/m ²	16.7±3.8	22.9±8.7	4.1±1.5	13.2±5.5
	Flowering plants/m ²	0.8±0.6	0.2±0.2	0.2±0.2	0.4±0.4
	Biomass (dry g/m ²)	19.0±4.8	9.7±3.6	1.6±1.3	4.9±2.3
Tyria density/m ²	6.4±3.4	5.1±2.6	0.5±0.4	0.4±0.3	

Study design	Timeseries		30pts
Baseline comparison	The soil type and size of each site was unreported		4pts
Intra treatment variation	All factors reported and suitably similar		4pts
Measurement of intervention and co-interventions	The number of <i>Tyria</i> per site/per year were recorded.		2pts
Replication & parameter of abundance	Unknown level of replication – transects were followed however number of quadrats were not listed. Suitable parameters of abundance measured.		1pt
Data Quality Score Total	Datasets A-D		41 pts
Other notes	Densities of 1 st year plants did not differ significantly, even though a downward trend occurred. Densities of flowering plants were significantly different ($P < 0.05$). Potential ragwort biomass was reduced in 4 year.		

Study 12	Pemberton, R. W. & Turner, C. E. (1990)			
Methods	A timeseries of three sites studying the impact that larvae of <i>Tyria jacobaeae</i> & <i>Longitarsus jacobaeae</i> have on common ragwort.			
Population	Size of experimental area:	Varies – see description below		
	Habitat:	Varies – see description below		
	Location:	3 sites near Fort Bragg, California, U.S.A.		
	Altitude:	Unreported		
	Plant age at time of treatment:	All ages		
	Soil type:	Unreported		
Weather	No weather was reported within the study.			
Intervention & Comparator	<p>Data from the following three sites were extracted:</p> <p>Site 1 – Foresti Ranch – ca. 4 ha of pasture of moist prairie on river bottomland (Ten Mile River); lightly grazed. <i>Longitarsus</i> ≥ 10/plant.</p> <p>Site 2 – Todd Point – ca. 30 ha coastal prairie; old pasture no longer grazed; vegetation now dense 1m high.</p> <p>Site 3 – Smith Ranch – ca. 1 ha pasture; heavily grazed; <i>Longitarsus</i> ≤ 10/plant. Sampling was undertaken in the Autumn measuring ragwort plants/m².</p>			
Outcomes		Foresti Ranch	Todd Point	Smith Ranch
	1966		<i>(Tyria only)</i> 53.3	
	1968			
	1969	<i>(Tyria already present – Longitarsus introduced)</i> 15.3		<i>(Tyria already present)</i>
	1972		71.1 <i>(Longitarsus introduced)</i>	<i>(Longitarsus introduced)</i>
	1973		39.5	11.7
	1974		6.9	7.4
	1975	0.0	0.6	0.5
	1976		0.6	0.2
	1987	0.0	0.0	0.18
Study design	Timeseries			30pts
Baseline comparison	The soil type and altitude of each site was unreported			4pts
Intra treatment variation	All factors are similar – expect altitude (unreported)			3pts
Measurement of intervention and co-interventions	The number of <i>Tyria</i> & <i>Longitarsus</i> per site/per m ² were recorded. Co-interventions were listed for each of the sites – namely grazing.			2pts
Replication & parameter of abundance	Suitable parameters of abundance measured, 200 quadrats taken for the mean ragwort plants/m ² however sd/se not reported.			1pt
Data Quality Score Total	Dataset A-C		40 pts	
Other notes				

Study 13	Schmidl, L. (1972)				
Methods	A control trial of the response of ragwort to defoliation of <i>Tyria jacobaeae</i> .				
Population	Size of experimental area:	Unreported			
	Habitat:	Abandoned dairy farm – overgrown pasture			
	Location:	Victoria, Australia			
	Altitude:	Unreported			
	Plant age at time of treatment:	All ages – natural population			
	Soil type:	Unknown			
Weather	Mean rainfall for the area is 1097mm, uniform throughout the year, averaging 60-80mm/month in summer & 100-110mm/month in winter & early spring. The hottest months are Jan & Feb but seldom exceed 39°C.				
Intervention & Comparator	The site was a dairy farm until abandoned in the 1930s. Area was overgrown with species of <i>Acacia</i> , <i>Cassinia</i> & <i>Rubus</i> with dominant grasses <i>Agrostis tenuis</i> , <i>Anthoxanthum odoratum</i> & <i>Holcus lanatus</i> . Plant characteristics were assessed in April/May, (4 month follow-up). n = 1 for level of replication.				
Outcomes		Dry weight (g)	Height (cm)	No. of capitula	Seed germination (%)
	Treatment	11	76	143	52
	Control	26	89	393	84
	Reduction (%)	58	15	64	38
Study design	Control Trial				60pts
Baseline comparison	The size of the experimental area; the altitude & soil type are unknown				3pts
Intra treatment variation	Again only altitude is unknown				3pts
Measurement of intervention and co-interventions	All parameters are known and similar. No co-interventions are reported.				2pts
Replication & parameter of abundance	Unknown level of replication. However parameters measured are sensible				1pt
Data Quality Score Total					69pts
Other notes	Feeding tests have shown that 15-20 larvae were sufficient to cause almost complete defoliation of small plants. 51% of ragwort plants attacked by a dense population of <i>Tyria</i> formed fresh growth and all regenerated plants produced a second crop of flowers and seeds. At the time of the assessment 72% of the recovered plants produced axial shoots, with an mean of 6 shoots/plant. 100% of the plants had signs of crown regeneration. Ragwort is a serious problem in the high-rainfall, high fertility areas of southern Victoria, Australia.				

Study 14	Windig, J. J. (1993)		
Methods	A timeseries assessing the effect of ragwort mortality via the investigation of two sites within the Dutch dunes.		
Population	Size of experimental area:	Two sites: Site 1 – 10x90m, Site 2 – unknown With sites 102 permanent 2x2m plots monitored.	
	Habitat:	Sand dunes/dune grasslands	
	Location:	Meijndel, Netherlands (52°08'N, 4°22'E)	
	Altitude:	<50m (Coastal Dune Lowlands)	
	Plant age at time of treatment:	Mixed – naturally	
	Soil type:	Sand based soils – mainly free draining.	
Weather	No detailed reporting of weather conditions was undertaken within the study. However frost damage was reported on plants.		
Intervention & Comparator	<p>Site 1 – dataset A Measuring 10x90 (900m²), positioned in a shady part of the dunes. Transition from open sands with minimal vegetation to dense vegetation consisting mostly of grasses. All <i>S. jacobaea</i> plants were marked. The rate of larval herbivory was estimated by sampling each month from January-June 1987. Ten randomly selected plants of various sizes were dug up and dissected to estimate the number of larvae per plant.</p> <p>Site 2 – dataset B An open area of dunes. The rate of herbivory was established as above. Plants on this site were at least double the size of site 1, with twice as many “shot-holes” of adult herbivory.</p> <p>The disappearance of above ground plant parts were checked every month for all plants in both sites, as was done for the regrowth. Also present on site were natural populations of <i>Tyria jacobaeae</i>. The results presented as percentage mortality – measured as a timeseries since January 1st 1987.</p>		
Outcomes		Site 1	Site 2
	Start	0	0
	End	60	80
	Longitarsus larvae /per plant	4.6	46.7
Study design	Timeseries		30pts
Baseline comparison	The altitude of each site was unreported		5pts
Intra treatment variation	Again only altitude is unknown		3pts
Measurement of intervention and co-interventions	Longitarsus numbers given, however <i>Tyria</i> numbers were unknown. No other co-interventions or activities to confound results.		2pts
Replication & parameter of abundance	Parameter measured suitable, however unknown replication.		1pt
Data Quality Score Total	Both dataset A-B	41pts	
Other notes	Twice at both sites (90 & 210 days) mortality of the <i>S. jacobaea</i> plants nearly reached 100%, however later regrowth meant that not all the plants had died. Mortality increased on both sites until 120 days after 1 st Jan then at site 1 – (shady area) mortality was near constant at 60% of all <i>S. jacobaea</i> plants. However at site 2 – mortality continued to steadily rise until 150 days then steadied peaking at 180 days (80%) before becoming constant. It was shown that Light Intensity was an important factor for the		